arXiv:1701.05371v1 [math.PR] 19 Jan 2017

Analytic solution for preferential attachment probabilities scheme

Andrea Monsellato

January 20, 2017

Abstract

Preferential attachment probabilities scheme appear in the context of scale free random graphs [1],[2]. In this work we present preferential attachment probabilities scheme as a sequence of dependent Bernoulli random variables and we give an explicit expression of the transition probabilities.

1 Introduction

Preferential attachment probabilities schemes arise in the context of scale free random graphs, following the description used in [2]:

The preferential linking arises in this simple model not because of some special rule including a function of vertex degree as in [1],[3] but quite naturally. Indeed, in the model that we consider here, the probability that a vertex has a randomly chosen edge attached to it is equal to the ratio of the degree k of the vertex and the total number of edges, $2t-1$.

we want to mimic this dynamic using a sequence of dependent Bernoulli random variables.

Let $\{Y_m(n)\}_{n>0}$ a sequence of r.v. such that $Y_m(n) \sim Ber\left(\frac{\sum_{i=m}^{n-1} Y_m(i)}{2n-1}\right)$, with $Y_m(1) = 1$. The sequence $\{Y_m(n)\}_{n>0}$, represent the trails to attach a new arcs at node m when a new node n is generated and jointed to the network, obviously $n > m$. The degree of the node m will be $X_m(n) = \sum_{i=m}^{n} Y_m(i)$.

We recover transition probabilities using elementary probability arguments, then

$$\begin{aligned}
P(X_m(n)=k) =& P(X_m(n)=k|X_m(n-1)=k-1)P(X_m(n-1)=k-1)+\\
&+P(X_m(n)=k|X_m(n-1)=k)P(X_m(n-1)=k)=\\
\\
&=P(X_m(n-1)+Y_m(n)=k|X_m(n-1)=k-1)P(X_m(n-1)=k-1)+\\
&+P(X_m(n-1)+Y_m(n)=k|X_m(n-1)=k)P(X_m(n-1)=k)=\\
\\
&=P(Y_m(n)=1|X_m(n-1)=k-1)P(X_m(n-1)=k-1)+\\
&+P(Y_m(n)=0|X_m(n-1)=k)P(X_m(n-1)=k)=\\
\\
&=\frac{k-1}{2n-1}P(X_m(n-1)=k-1)+\frac{2n-1-k}{2n-1}P(X_m(n-1)=k)
\end{aligned}$$

Now set $p^m_{n,k}=P(X_m(n)=k)$, we have that

$$p^m_{n,k}=p^m_{n-1,k-1}\frac{k-1}{2n-1}+p^m_{n-1,k}\frac{2n-1-k}{2n-1},\quad 0\le m<n,\quad 0\le k\le n-m \tag{1}$$

2 Recovering explicit expression of transition probabilities

Choosing $m=0$ in (1), i.e. we consider the probabilities transition equations for the first node, we have

$$\begin{cases} p_{1,1}=1,\quad p_{n,k}=0,\quad \forall k>n\\ p_{n,k}=\frac{k-1}{2n-1}p_{n-1,k-1}+\frac{2n-1-k}{2n-1}p_{n-1,k},\quad \forall k\le n \end{cases} \tag{2}$$

Observe that (1) is time invariant for all nodes scaling properly the quantity $n-m$.

For (2) holds that:

$$p_{n,1}=\frac{2(n-1)}{2n-1}p_{n-1,1}\quad \forall n\ge 2$$

i.e.

$$p_{n,1}=\frac{2^{n-1}(n-1)!}{\prod_{i=1}^{n}(2i-1)}\quad \forall n\ge 1 \tag{3}$$

also

$$p_{n,n}=\frac{n-1}{2n-1}p_{n-1,n-1}\quad \forall n\ge 2$$

i.e.

$$p_{n,n}=\frac{(n-1)!}{\prod_{i=1}^{n}(2i-1)}\quad \forall n\ge 1$$

Let

$$a_{n,k} = \frac{2^{n-k}\prod_{i=1}^{n}(2i-1)}{(n-1)!}p_{n,k} \tag{4}$$

from (2),(3) and (4) we have that

$$\begin{cases} a_{n,1} = 2^{2(n-1)}, \quad a_{n,n} = 1, \quad a_{n,k} = 0 \quad \forall k > n \\ a_{n,k} = \frac{1}{n-1}\left[(k-1)a_{n-1,k-1} + 2(2n-1-k)a_{n-1,k}\right] \quad \forall k = 1, ..., n-1 \quad n > 1 \end{cases} \tag{5}$$

Now we give an analytic solution of (5).

Theorem 1. *Let $\{a_{n,k}\}$ as in (5), then*

$$\begin{cases} a_{n,n} = 1 \\ a_{n,k} = 2^{2(n-k)}\left[1 + (k-1)\sum_{j=0}^{n-k-1}\frac{2^{-2(j+1)}(k+2j)!}{(j+1)!(k+j)!}\right] \quad \forall k = 1, ..., n-1 \quad n > 1 \end{cases} \tag{6}$$

Proof

If $k = 1$ obvious.

If $k = n-1$ and $\forall n \geq 2$, then (5) will be

$$a_{n,n-1} = \frac{1}{n-1}\left[(n-2)a_{n-1,n-2} + 2na_{n-1,n-1}\right] \tag{7}$$

now substituting (6) in (7) we have that

$$2^2\left[1 + (n-2)\sum_{j=0}^{0}\frac{2^{-2(j+1)}(n-1+2j)!}{(j+1)!(n-1+j)!}\right] =$$

$$= \frac{n-2}{n-1}2^{2(n-1-n+2)}\left[1 + (n-3)\sum_{j=0}^{n-1-n+2-1}\frac{2^{-2(j+1)}(n-2+2j)!}{(j+1)!(n-2+j)!}\right] + \frac{2n}{n-1}$$

$$2^2 + (n-2) = \frac{n-2}{n-1}2^2\left[1 + (n-3)2^{-2}\right] + \frac{2n}{n-1}$$

$$2^2(n-1) - 2n = 2n - 4$$

$$-4 = -4$$

Let $k = 2, ..., n-2$ substituting we have that

$$(n-1)\left[1+(k-1)\sum_{j=0}^{n-k-1}\frac{2^{-2(j+1)}(k+2j)!}{(j+1)!(k+j)!}\right] = (k-1)\left[1+(k-2)\sum_{j=0}^{n-k-1}\frac{2^{-2(j+1)}(k-1+2j)!}{(j+1)!(k-1+j)!}\right]$$

$$(n-1)\left[1+(k-1)\sum_{j=0}^{n-k-1}\frac{2^{-2(j+1)}(k+2j)!}{(j+1)!(k+j)!}\right] = \frac{1}{2}(2n-1-k)\left[1+(k-1)\sum_{j=0}^{n-k-2}\frac{2^{-2(j+1)}(k+2j)!}{(j+1)!(k+j)!}\right]$$

$$(n-1)\left[1+(k-1)\sum_{j=0}^{n-k-1}\frac{2^{-2(j+1)}(k+2j)!}{(j+1)!(k+j)!}\right] = \left[(n-1)-\frac{1}{2}(k-1)\right]\left[1+(k-1)\sum_{j=0}^{n-k-2}\frac{2^{-2(j+1)}(k+2j)!}{(j+1)!(k+j)!}\right] \tag{8}$$

we recover that (8) is equivalent to

$$\sum_{j=0}^{n-k-2}\frac{2^{-2(j+1)}}{(j+1)!}\left[2(k-2)\frac{(k-1+2j)!}{(k-1+j)!}-(k-1)\frac{(k+2j)!}{(k+j)!}\right] = 2^{-2(n-k-1)}\frac{(2n-k-3)!}{(n-k-1)!(n-2)!}-1 \tag{9}$$

Now let $n = k+2+r$, $r \geq 0$ the equality (9) become

$$\sum_{j=0}^{r}\frac{2^{-2(j+1)}}{(j+1)!}\left[2(k-2)\frac{(k-1+2j)!}{(k-1+j)!}-(k-1)\frac{(k+2j)!}{(k+j)!}\right] = 2^{-2(r+1)}\frac{(k+1+2r)!}{(r+1)!(k+r)!}-1 \tag{10}$$

By induction we prove equality (10):

For $r = 0$ we have

$$\frac{2^{-2}}{1!}[2(k-2)-(k-1)] = \frac{2^{-2}(k+1)!}{1!k!}-1$$

i.e. $2^{-2}(k-3) = 2^{-2}(k-3)$.

By induction hypothesis we prove equality (10) for $r+1$, supposing it holds for generic r, then:

$$2^{-2(r+1)}\frac{(k+1+2r)!}{(r+1)!(k+r)!}+\frac{2^{-2(r+2)}}{(r+2)!}\left[2(k-2)\frac{(k+1+2r)!}{(k+r)!}-(k-1)\frac{(k+2+2r)!}{(k+1+r)!}\right] = 2^{-2(r+2)}\frac{(k+3+2r)!}{(r+2)!(k+r+1)!}$$

$$2^{2(r+2)}\frac{(k+1+2r)!}{(k+r)!}+2(k-2)\frac{(k+1+2r)!}{(k+r)!}-(k-1)\frac{(k+2+2r)!}{(k+1+r)!} = \frac{(k+3+2r)!}{(k+r+1)!}$$

i.e.

$$2(r+2)(k+r+1)+2(k-2)(k+r+1)-(k-1)(k+2+2r)=(k+3+2r)(k+2+2r)$$

from obvious arrangement w.r.t. k the thesis follow.

Now consider that $\prod_{i=1}^{n}(2i-1)=\frac{(2n)!}{2^n n!}$, for $k=1,...,n-1$ holds that

$$p_{n,k}=\frac{(n-1)!}{2^{n-k}\prod_{i=1}^{n}(2i-1)}a_{n,k}=\frac{(n-1)!}{2^{n-k}(2n)!}2^n n!a_{n,k}$$

i.e.

$$p_{n,k}=\frac{n!(n-1)!}{(2n)!}2^{2n-k}\left[1+(k-1)\sum_{j=0}^{n-k-1}\frac{2^{-2(j+1)}}{(j+1)!}\frac{(k+2j)!}{(k+j)!}\right]$$

References

[1] R.,Albert, A.L., Barabsi, Statistical mechanics of complex networks, Rev. Mod. Phys, 2002

[2] S.N., Dorogovtsev, J.F.F., Mendes, Evolution of networks, Adv. Phys., 2002, 51, 1079

[3] S.N., Dorogovtsev, J.F.F., Mendes, A.N., Samukhin, Structure of Growing Networks with Preferential Linking, Physical Review Letters, 2000, 21